

Quotes: DPT vaccines

Quotes

See: [Diphtheria vaccine quotes](#) [Whooping Cough quotes](#) [Tetanus quotes](#) [Diphtheria quotes](#)

"Several factors indicate that mass immunisation with pertussis and other non-Hib vaccines may have been responsible for the unprecedented epidemics of invasive bacterial infections such as [Hib](#), during the 1970's and 1980's." [Vaccine Safety Manual by Neil Z. Miller. \(p.315\)](#)

[\[30 Jan 2008\] U of M researcher links asthma, early vaccinations](#)

Kozyrskyj found nearly 14 per cent of the children who received their first shot of diphtheria, pertussis, tetanus vaccine at two months of age developed asthma — compared with only 5.9 per cent of children who were vaccinated more than four months after the scheduled date.

[\[Jan 2008\] The Federal Government recently conceded a real vaccine-autism lawsuit in a real court](#)

"The Federal Government recently conceded a [real vaccine-autism lawsuit](#) in a real court and will soon pay a real (taxpayer-funded) settlement to a real American family and a very real child with autism.....If I were the AAP, or ABC for that matter, I would feel downright silly stating that "no scientific link exists," so soon after the [Journal of Child Neurology](#) published a study titled, "Blood Levels of Mercury Are Related to Diagnosis of Autism: A Reanalysis of an Important Data Set." I would also worry about parental reaction to learning that researchers had done due diligence and reanalyzed data from a prior, hugely influential study that (erroneously) found zero connection between mercury levels and autism.....

Another study, freshly out of Harvard, likewise shows a potential link between mercury and the autopsied brains of young people with autism. The [American Journal of Biochemistry and Biotechnology](#) reports that a marker for oxidative stress was 68.9% higher in autistic brain tissue than controls (a statistically significant result), while mercury levels were 68.2% higher.

In the article, "Delay in DPT vaccination is associated with a reduced risk of childhood asthma," Anita Kozyrskyj, an asthma researcher at the University of Manitoba, and other scientists combed the medical records of 14,000 children born in Manitoba in 1995 (when many Canadian shots still contained mercury, by the way). They found that children who received the DPT (diphtheria, pertussis and tetanus) vaccine at two months of age were 2.63 times more likely to develop asthma (at a rate of 13.9%) than children who were not given the shot until after four months of age (5.9%)."

Searching for children who had not been exposed to mercury in vaccines -- the kind of population that scientists typically use as a "control" in experiments -- [Dan Olmsted](#) scoured the Amish of Lancaster County, Pennsylvania, who refuse to immunize their infants. Given the national rate of autism, Olmsted calculated that there should be 130 autistics among the Amish. He found only four. One had been exposed to high levels of mercury from a power plant. The other three -- including one child adopted from outside the Amish community -- had received their vaccines.--
--[Deadly Immunity By ROBERT F. KENNEDY JR.](#)

"When in Germany, vaccinations against whooping cough were suspended between 1975 and 1991, the declining trend in the number of deaths from the disease continued as before."--Dr [Buchwald](#) (The Decline of Tuberculosis despite "Protective" Vaccination by Dr. Gerhard Buchwald M.D. p135)

Bhubaneswar, India : Seventy-nine schoolchildren in Orissa have fallen ill, 11 of them critically, after they were given tetanus and diphtheria vaccination shots that are part of the state's immunisation programme. Officials said 47 students at the Kodakana primary school in Kendrapada district were given the tetanus toxoid shots and 32 were given the diphtheria toxoid vaccination on Aug 26 in the school premises by local paramedical staff. After three days, all the students complained of skin irritation, viral fever and vomiting. Eleven of them are critically ill and have been admitted to the local government hospital. One of them, a 10-year-old girl called Minirani Khatua, has blood oozing from her nose and mouth, said a district health official. [\[Media Sept 2004\] India News: 79 schoolchildren ill after vaccinations](#)

Recently, my VACCINATED DPT autistic child came down with PERTUSSIS (which he lived through just fine). While in the ER, I came across four other nurses with a child with autism, who in secret meetings to me between doctors and tests, revealed that they too felt their children were vaccine induced autistic. [\[Letter BMJ 19 July 2004\] Cathy Blanco](#)

[2004 jan] In conclusion, for those who have a decreased ability to excrete mercury, as has been demonstrated for several different genotypes, there can be little doubt that mercury concentrations once administered to children as part of the childhood routine vaccination schedule resulted in a significant number of children developing autism. This is especially true following a sudden increase in the amount of mercury administered, as occurred in the United States in the early 1990s when the amount of mercury administered to children in the first six months of life more than doubled as part of the routine childhood immunization schedule (i.e. from 75 micrograms of mercury from three DTP immunizations to 187.5 micrograms from three DTP, three Hib, and three hepatitis B immunizations).....It is also clear that if somehow, despite the over whelming evidence, the IOM determines, that either thimerosal did not cause

or that they are not sure that it caused the current epidemic of autism and other neurological disorders, that the IOM must demand the immediate expenditure of billions of dollars as part of an all out effort to immediately determine what is causing this epidemic before it totally destroys our society. [\[jan 2004\] A Review of the Relationship between Thimerosal and Autism. David A. Geier and Mark R. Geier, MD](#)

"How can the Tetanus vaccine induce immunity, when contracting the disease naturally does not give immunity?"--NVIC

An astonishing journal paper. 1 November, 1993. FASEB Journal, volume 7, pp.1381-1385. Authors—Stephan Dirnhofer et al. Dirnhofer is from the Institute for Biomedical Aging Research of the Austrian Academy of Sciences. A quote from the paper: "Our study provides insights into possible modes of action of the birth control vaccine promoted by the Task Force on Birth Control Vaccines of the WHO (World Health Organization)." A birth control vaccine? What? Yes. A vaccine whose purpose is to achieve non-pregnancy where it ordinarily could occur. Sterilization? This particular vaccine is apparently just one of several anti-fertility vaccines the Task Force is promoting. Yes. There is a Task Force on Birth Control Vaccines at WHO. This journal paper focuses on a hormone called human chorionic gonadotropin B (hCG). There is a heading in the paper (p.1382) called "Ability of antibodies to neutralize the biological activity of hCG." The authors are trying to discover whether a state of no-fertility can be achieved by blocking the normal activity of hCG. They state, "We conclude from our results that both the efficacy and safety of the WHO vaccine are not yet ensured."

Another journal paper. The British Medical Bulletin, volume 49,1993. "Contraceptive Vaccines" is the title of the paper. The authors—RJ Aitken et al. From the MRC Reproductive Biology Unit, University of Edinburgh, Edinburgh, UK.

"Three major approaches to contraceptive vaccine development are being pursued at the present time. The most advanced approach, which has already reached the stage of phase 2 clinical trials [human trials testing efficacy], involves the induction of immunity against human chorionic gonadotrophin (hCG). Vaccines are being engineered ... incorporating tetanus or diphtheria toxoid linked to a variety of hCG-based peptides... Clinical trials have revealed that such preparations are capable of stimulating the production of anti-hCG antibodies. However, the long-term consequences of such immunity in terms of safety or efficacy are, as yet, unknown...The authors are talking about creating an immune response against a female hormone....The authors state, "The fundamental principle behind this approach to contraceptive vaccine development is to prevent the maternal recognition of pregnancy by inducing a state of immunity against hCG, the hormone that signals the

presence of the embryo to the maternal endocrine system."..... [Rappoport, Jon](#) (Ownership of All Life p66)

"Capps R.B.et al (1955) stated DPT caused temporary liver dysfunction in infants similar to that caused by viral hepatitis. Anser S and Habig W (1990) showed DPT vaccine endotoxin significantly disrupts P-450, and other microsomal and cytosolic enzyme activities (which detoxify endotoxin) in mice. Since gastrointestinal E.coli is detoxified by the liver, any suppression of the RE (Reticulo endothelial) system by vaccines places a baby at risk of death by endotoxin effect. Rook, G.A.W (1997) details other ways vaccines disrupt the immune system."---[Hilary Butler](#)

"The whooping cough vaccine has never had much of an influence on the number of children dying from whooping cough. The dramatic fall in the number of deaths caused by the disease came well before the vaccine was widely available and was, historians agree, the result of improved public health measures and, indirectly, the use of antibiotics. The introduction of the vaccine really didn't make very much, if any, difference to the fall in the incidence of the disease. Even today (1988) thirty years after the introduction of the vaccine, whooping cough cases are still running at about 1,000 a week in Britain. Similarly, the figures show that the introduction of the vaccine had no effect on the number of children dying from whooping cough.My second point is that the whooping cough vaccine is neither very efficient nor is it safe..... the government (in the form of the DHSS) has consistently lied about the whooping cough vaccine, has distorted the truth and has deceived both the medical profession (for the majority of doctors and nurses who give these injections accept the recommendations made by the DHSS without question) and millions of parents. The DHSS may have saved itself a tidy sum in damages. But the cost to the nation's health has been enormous. And this, remember, is merely one more example of the way in which the truth has been distorted by those whom we trust to provide us with honest, accurate advice about medicine and health care. "---- [Dr Vernon Coleman MB](#)

"Did you know that the whooping cough germ, Bacillus pertussis, when injected into animals, has long been known to lead to the secretion of insulin? In 1979, at the Fourth International Symposium on Pertussis, held in Bethesda, Maryland, it was shown that this same result occurs in those who have received pertussis vaccine. In their publication, "Adverse Reactions after Pertussis Vaccination," Drs. W. Hennesen and U. Quast suggest, "It seemed of interest to examine these reactions in comparison with the hypoglycemia syndrome. . . There was a close relation between the two.' If your child has juvenile diabetes (a disease characterized by wide swings in blood sugar levels), ask your doctor if he has ever heard of this effect of whooping cough vaccine. Maybe it's time to investigate whether the pertussis vaccine has anything to do with the rapidly rising number of people with juvenile diabetes, adult diabetes, and

hypoglycemic all disorders of insulin metabolism."---[Dr Robert Mendelsohn MD](#) (the Peoples Doctor Vol 6 No10)

"At least since the 1970s, pertussis vaccine has been known in animal experiments to stimulate over-production of insulin by the pancreas followed by exhaustion and destruction of the "islets" with consequent under-production of insulin; in the first case the outcome is hypoglycemia, and in the latter it is diabetes. Physicians as early as 1949 called attention to low blood glucose in children who had severe reactions to the pertussis vaccine. In 1970, Margaret Pittman wrote: "the infant whose blood sugar level is influenced by food intake may be especially vulnerable to vaccine-induced hypglycemia...the vaccine induces hypoglycemia in mice and rabbits." Gordon Stewart wrote in 1977: "more than any other vaccine in common use, pertussis vaccine is known pharmacologically to provoke...hypoglycemia due to increase production of insulin." Two Dutch researchers, Hannik and Cohen, observed in 1978: "infants who show serious reactions following pertussis vaccination suffer from a failure to maintain glucose homeostasis." And two German researchers, Hennessen and Quast, found in 1979 that 59 out of 149 children who manifested adverse reactions to the pertussis vaccine developed symptoms of hypoglycemia."---[Harris Coulter](#)

"At least half of all U.S children have had otitis media by their first birthday. By age 6 90% have had them. This condition accounts for 26 million visits to physicians every year. In addition, about 1 million children have tubes inserted in their ears every year, at a cost of \$1000/operation. Thus \$1 billion is spent each year on this operation. Just imagine what it means if this is all, or mostly all, caused by the pertussis vaccine. This particular "glue ear" type of otitis was not known in American medical practice before the late 1940's or early 1950's -- in other words, the time when the pertussis vaccine was being introduced."--[Harris Coulter Ph.D.](#)

"It now transpires that during the Clinical Trials children did in fact suffer severe and irreversible brain damage as a result of the Pertussis Vaccine and accepted as such by Medical Experts appointed by the Department of Health.....It is now an established fact that the present figure now stands at over 900 which is an absolute disgrace in a civilised country."--[M Wain http://www.whale.to/vaccines/wain.html](http://www.whale.to/vaccines/wain.html)

"Killed bacteria may release toxins as their cell bodies break up. An example is pertussis vaccine, which contains at least one substance that can be poisonous to brain cells. When the toxin injures the brain, this occurs anywhere from a few hours to a few days after the vaccination."--Dr Kingsbourne
MD<http://www.whale.to/vaccines/kingsbourne3.html>

"In a year-long investigation of the Vaccine Adverse Reaction Reporting System (VAERS) operated by the Food and Drug Administration, NVIC/DPT analyzed VAERS computer discs used by the FDA to store data on reports of deaths and injuries following DPT vaccination. A total of 54,072 reports of adverse events following vaccination were listed in a 39-month period from July 1990 to November 1993 with 12,504 reports being associated with DPT vaccine, including 471 deaths." [Campaign Against Fraudulent Medical Research \(CAFMR\)](#)

"For decades, American pharmaceutical companies have known how to produce the safer DPT vaccine but decided not to bring it to market because it would increase production costs and lower the drug's 50% or higher profit margins."--[Money Magazine](#)

"When the link between the use of unsafe, mercury-laden vaccine and autism, ADHD, asthma, allergies and diabetes becomes undeniable, mainstream medicine will be sporting a huge, self-inflicted and well-deserved black eye. Then will come the billion-dollar awards, by enraged juries, to the children and their families. I can't wait."--[Dr Rimland MD](#)

"The role of vaccines, particularly the whole-cell pertussis (whooping cough) vaccine can be understood when it is realised that this vaccine contains a variable and uncontrollable amount of endotoxin that is injected and absorbed, unaltered, into the blood. It does not even go first to the liver where attempts to detoxify it could be made. If an infant happens to be particularly sensitive to endotoxin when the vaccine is injected, brain damage or death can result....It should be now apparent that any infant with gastrointestinal problems - abnormal organisms, intestinal parasites, loose bowel motions resulting from the use of antibiotics, and malabsorption of food (including lactose intolerance) - is liable, when further stressed, to produce endotoxin and this can end in a SIDS....If the Vitamin C status of an infant is borderline, the administration of a vaccine, particularly (but not only) pertussis vaccine, can result in endotoxaemia. This results in a severe reaction to the vaccine, a tremendous increase in the need for Vitamin C, and the precipitation of some of the signs and/or symptoms of acute scurvy. The onset of this may be so rapid that the classical signs of scurvy may be absent. Sudden death, sudden unconsciousness, sudden shock or sudden spontaneous bruising and haemorrhage (including brain and retinal haemorrhages) may occur. Haemorrhage and bruising in such cases can be wrongly attributed to the 'battered baby syndrome'."---[Dr Kalokerinos MD](#) (Medical Pioneer of the 20th century p186)

"The seldom publicized history of the pertussis vaccine in Sweden, gives an entirely different point of view from that of the US public health service. Sweden banned the pertussis vaccine in 1979, and yet Sweden now has the second lowest infant mortality

rate in the world, whereas the United States ranks a very poor 20th or lower.....a report ([Trollfors B](#). Bordetella pertussis whole cell vaccines--efficacy and toxicity. Acta Paediatr Scand. 1984 Jul;73(4):417-25. Review. PMID: 6380211; UI: 84276893.) in 1984 stated that the pertussis mortality was generally very low in industrialized countries and there was no difference in severity and incidence of pertussis between countries with high, low and zero immunizations rates."--[Harold Buttram](#)

A study revealed that 1 in 175 children who completed the full DPT suffered severe reactions ("[Nature and Rates of Adverse Reactions Associated with DPT and DT Immunizations in Infants and Children](#)" [Paediatrics, Nov. 1981, Vol.68, No.5]) and a Dr.'s report for attorneys which found that 1 in 300 DPT immunizations resulted in seizures (The Fresno Bee, Community Relations, DPT Report, Dec 5, 1984).

"Every day new parents are ringing us. They all have the same tragic story. Healthy baby, child, teenager, usually a boy, given the DPT (diphtheria, pertussis and tetanus) or DT (diphtheria and tetanus), MMR or MMR booster followed by a sudden fall or slow, but steady decline into autism or other spectrums disorder."--[The Hope Project](#)

"Delay of DPT immunization until 2 years of age in Japan has resulted in a dramatic decline in adverse side effects. In the period of 1970-1974, when DPT vaccination was begun at 3 to 5 months of age, the Japanese national compensation system paid out claims for 57 permanent severe damage vaccine cases, and 37 deaths. During the ensuing six year period 1975-1980, when DPT injections were delayed to 24 months of age, severe reactions from the vaccine were reduced to a total of eight with three deaths. This represents an 85 to 90 percent reduction in severe cases of damage and death. (Ref21)."-[Dr Obomsawin MD](#)

"Classen's data and other published data indicates the following vaccines are associated with an increased risk of diabetes (increased risk): hepatitis B (50%), hemophilus (25%), tetanus (20%), diphtheria (9%), pertussis (25%), mumps- rubella (23%). These findings are supported by a case control study performed in Europe. The cumulative effect of all these vaccines on diabetes is tremendous."--[PRNewswire](#)

"Dr. Kevin Geraghty, Bay Area Physicians for the Study of Pertussis Vaccines: "The UCLA study showed one in 600 kids have shock like episodes. That is not controversial.....I'm convinced in my heart of hearts that 100 American infants a year are dying and another 250 are brain-damaged from DPT. I would be prepared to clinically defend that to scientists." Geraghty believes as many as one in 15,000 children suffer brain damage and as many as one in 35,000 die following DPT vaccination."---[Fresno Bee, DPT report 1984](#).

"My own view, based upon some years of observation and experience, is quite firm. I supported the use of the vaccine in 1951 and subsequently with very little hesitation until about 1972, and gave pertussis vaccine between 1951 and 1956 to each of my four children. I would not dream of doing so again because it has become clear to me not only that the vaccine is incompletely protective, but also that the side-effects which I thought to be temporary are in fact dangerous, unpredictably so. There is no doubt in my mind that in the UK alone some hundreds, if not thousands, of well infants have suffered irreparable brain damage needlessly and that their lives and those of their parents have been wrecked in consequence."--[Prof Gordon Stewart 1980](#)